




REGIONAL CHARACTERIZATION

Regional characterization activities increase understanding of the magnitude, distribution, and variability of major stationary CO₂ sources and potential CO₂ geologic storage sites. Ongoing regional characterization in the PCOR Partnership region supports CO₂ storage project development through the acquisition and analysis of subsurface data to help scientists, engineers, and project developers understand the relevant properties and characteristics of the subsurface environment. These characterization efforts are a necessary step in CCUS project development where the ideal pairings of industrial facilities that can capture CO₂ and suitable geologic storage targets can be identified.




DISTRIBUTION OF LARGE STATIONARY CO₂ SOURCES




CO₂ SOURCES

The PCOR Partnership has identified, quantified, and categorized 512 stationary sources in the region that have an annual output of greater than 100,000 tonnes of CO₂. These stationary sources have a combined annual CO₂ output of over 509 Mt. Although not a target source of CO₂ for geologic storage, the transportation sector in the U.S. portion of the PCOR Partnership region contributes nearly 242 million additional tonnes of CO₂ to the atmosphere every year.^{15,16,34-38}

The annual output from the various large stationary sources ranges from 100,000 tonnes for industrial and agricultural processing facilities that make up the majority of the sources in the region to over 14 Mt for the largest coal-fired electric generation facility. Fortunately, many of the large point sources are located in areas that are favorable for CO₂ storage because of their concurrence with deep sedimentary basins, such as those areas in Alberta, North Dakota, Montana, and Wyoming.



MAJOR REGIONAL SEDIMENTARY BASINS



Favorable geologic conditions for CO₂ storage exist in sedimentary basins found across the PCOR Partnership region.


Sedimentary Basin
(nominal extent)

CO₂ STORAGE OPPORTUNITIES


Sedimentary basins are large regional depressions in Earth's crust. These depressions accumulate a considerable thickness of sediment that can cause further subsidence and allow for even more sediments to accumulate. As the sediments are buried, they are subjected to compaction from increasing pressure and then begin the process of lithification (changing to rock). Sedimentary basins vary in configuration from bowl-shaped to elongated troughs. If organic-rich sedimentary rocks occur in combination with appropriate depth, temperature, and duration of burial, hydrocarbon generation can occur within the sedimentary basin. The rich set of options for the safe, long-

term geologic storage of CO₂ in the PCOR Partnership region is found in the deep portions of the extensive sedimentary basins of this region.

Oil and gas reservoirs and deep saline formations are the two primary CO₂ storage options found within sedimentary basins. These storage formations are commonly situated vertically one above another and separated by sealing formations, an arrangement referred to as stacked storage. Stacked storage offers the potential to store the same total volume of CO₂ but with a smaller geographic footprint.



EOR POTENTIAL



CO₂ STORAGE IN OIL FIELDS



Although oil was discovered in the PCOR Partnership Region in the late 1800s, significant development and exploration did not begin until the late 1920s. The body of knowledge gained in the nearly 90 years of exploration and production of hydrocarbons in this region is a significant step toward understanding the mechanisms for secure storage of significant amounts of CO₂. Today, oil is drawn from the many oil fields in the PCOR Partnership region from depths ranging from as little as 60 m to approximately 8000 m below ground level.

While the use of CO₂ in conventional reservoirs is a widely applied practice, the use for EOR in unconventional (or tight) oil reservoirs like the Bakken petroleum system (Bakken and Three Forks Formations) is a relatively new concept. Initial laboratory and field testing offers promising results that CO₂ for EOR in the Bakken may be a viable option. Current research is evaluating approaches to use CO₂ to improve Bakken oil production through field-scale injection testing. If proven viable, CO₂ EOR in unconventional reservoirs presents an opportunity for tremendous volumes of CO₂ storage and increases in oil production.


The region has over 3.7 Gt of CO₂ storage potential in conventional oil fields and 10.2 billion stb of incremental oil potential.



CO₂ STORAGE IN SALINE FORMATIONS

Characterization efforts of deep saline formations (DSFs) in the PCOR Partnership region indicate the potential to store over 330 Gt of CO₂. These DSFs often occur in stacked storage situations with EOR opportunities or other DSFs. The extent of the saline formations identified for storage is constrained by depth (to ensure the injected CO₂ remains in a dense liquid state) and by salinity (to protect groundwater sources).

Storage Potential in Deep Saline Formations million tonnes/100 square kilometers



CO₂ STORAGE IN COAL

The PCOR Partnership region is home to significant coal resources. Much of this vast resource is used to generate electricity at coal-fired power plants in the region and beyond. However, a significant portion of this resource lies at depths that are not economically recoverable.

The evaluation of three major coal horizons in the PCOR Partnership region identified nearly 7.3 Gt of CO₂ storage resource. However, because most of the coal resource in the PCOR Partnership region is positioned in the freshwater horizons of the subsurface, no current activities or potential CCUS projects are looking to use coal as a CO₂ storage target.



Major Coal Basin Regions of North America