

PCOR Partnership
ATLAS
2nd Edition



PCOR Partnership Atlas

2nd Edition

**Wesley D. Peck, Barry W. Botnen, Lisa S. Botnen, Daniel J. Daly, John A. Harju, Melanie D. Jensen, Erin M. O’Leary,
Steven A. Smith, James A. Sorensen, Edward N. Steadman, Stephanie L. Wolfe**

**Energy & Environmental Research Center
Grand Forks, North Dakota**

**Darin R. Damiani
John T. Litynski
U.S. Department of Energy**

**David W. Fischer
Fischer Oil and Gas, Inc.**

Prepared for the

**U.S. Department of Energy
National Energy Technology Laboratory
and the PCOR Partnership**

Published by the

**Energy & Environmental Research Center
2007**

The Plains CO₂ Reduction (PCOR) Partnership is a group of public and private stakeholders working together to better understand the technical and economic feasibility of sequestering CO₂ emissions from stationary sources in the central interior of North America. The PCOR Partnership is managed by the Energy & Environmental Research Center (EERC) at the University of North Dakota and is one of seven regional partnerships under the U.S. Department of Energy’s Regional Carbon Sequestration Partnership Program.

NOTICE

This document was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Department of Energy, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, expressed or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights.

Printed in the United States of America and
Available from:

Energy & Environmental Research Center
Grand Forks, ND 58203

This product was prepared by the Energy & Environmental Research Center (EERC), an agency of the University of North Dakota, as an account of work sponsored by the United States Department of Energy. Because of the research nature of the work performed, neither the EERC nor any of its employees makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement or recommendation by the EERC.

© 2007 University of North Dakota
Energy & Environmental Research Center

Permission is granted to copy and distribute information for noncommercial use, as long as the content remains unaltered and credit is given to the EERC. To commercially publish any of the materials included in this publication, please contact the EERC to obtain written permission. Write Derek Walters, EERC Communications Manager, 15 North 23rd Street, Stop 9018, Grand Forks, ND 58202-9018.

Acknowledgments

This was made possible through the contributions and efforts of numerous groups from throughout the United States and Canada. We gratefully acknowledge the individuals who provided the backbone information upon which this atlas is based, specifically, Stefan Bachu, Dean Bangsund, Randy Burke, Chip Euliss, Jim Evans, Robert Gleason, Lynn Helms, W.A. Jackson, Dick Kempka, Julie LeFever, Richard LeFever, Larry Leistriz, Claudia Miller, Ed Murphy, Harvey Ness, Ron Ness, Myria Perry, Randy Renner, Bill Reynen, Anne-Marie Thompson, Steve Whittaker, and Malcolm Wilson. We also acknowledge the PCOR Partnership partners for their valuable efforts in providing much of the information used for these assessments and for cooperating with us in producing a regional portfolio for public use.

We extend our appreciation to the various federal, state, and private organizations and university groups for their cooperation in our search for data.

A number of individuals at the Energy & Environmental Research Center provided valuable support for this effort. We are grateful to Earl Battle and Paul Gronhovd for their work in producing many of the graphics used in this document and to Joyce Riske for editing the atlas.

Table of Contents

Introduction	1
A Change Is in the Air	3
Greenhouse Effect	5
What Is CO ₂ ?	7
Carbon Management.....	9
Geologic Sequestration	11
Terrestrial Sequestration.....	13
DOE's Carbon Sequestration Regional Partnerships	15
The PCOR Partnership.....	17
PCOR Partnership Partners	19
Anthropogenic Sources	21
CO ₂ Sources	23
CO ₂ Source Types	25
Geologic Framework.....	27
Sedimentary Basins	29
Oil and Gas Fields.....	31
Enhanced Oil Recovery	33
Saline Formations	35
Sequestration in Coal	37
Terrestrial Sequestration.....	39
Prairie Pothole Region	41
Field Validation Test Sites	43
Carbon Capture and Separation	45
Regulatory and Safety Aspects of Carbon Capture and Storage.....	47
The PCOR Partnership Decision Support System	49
Keeping the Lights On	51
Education and Outreach – CO ₂ Sequestration	53
References and Photo/Image Credits	54



Introduction

Global climate change is considered to be one of the most pressing environmental concerns of our time. This is due in part to the potential magnitude of the changes it could cause and also to the immense economic, technological, and lifestyle changes that may be necessary in order to respond to it. Although uncertainty still clouds the science of climate change, there is strong indication that we may have to significantly reduce anthropogenic greenhouse gas (GHG) emissions. Carbon sequestration offers a promising set of technologies through which carbon dioxide (CO₂) and potentially other GHGs can be stored for long periods of time in sinks represented by biologic materials, geologic formations and, possibly, other places such as oceans. Within central North America, the Plains CO₂ Reduction (PCOR) Partnership is investigating sequestration technologies in order to provide a safe, effective, and efficient means of managing the carbon dioxide emissions across the center of the continent.

The regional characterization activities conducted under the initial effort of the PCOR Partnership confirmed that while there are numerous large stationary CO₂ sources, the region also has a variety of sinks that represent a tremendous capacity for CO₂ sequestration. The varying sources reflect the geographic and socioeconomic diversity of the region. In the upper Mississippi River Valley and along the shores of the Great Lakes Michigan and Superior, large coal-fired electrical generators power the manufacturing plants and breweries of St. Louis,


Minneapolis, and Milwaukee. To the west, the prairies and badlands of the north-central United States and central Canada are home to coal-fired power plants, natural gas processing plants, ethanol plants, and refineries that further fuel the industrial and domestic needs of cities throughout North America.

The PCOR Partnership region is rich in agricultural lands, forests, and wetlands that hold tremendous potential for terrestrial sequestration. The Prairie Pothole Region, which stretches from northwestern Iowa, across the Dakotas, and into Saskatchewan and Alberta, holds promise as an area that can be transformed into a significant terrestrial CO₂ sink. Deep beneath the surface of the region lay geologic formations that hold tremendous potential to store CO₂. Oil fields well suited for sequestering CO₂ can be found in roughly half the region, while formations of limestone, sandstone, and coal suitable for CO₂ storage exist in basins that, in some cases, extend over thousands of square miles. In many cases, large sources in the region are proximally located to large-capacity sinks, and in some cases, key infrastructure is already in place.

This atlas provides an introduction into the concept of global climate change and a regional profile of CO₂ sources and potential sinks across nearly 1.4 million square miles of the PCOR Partnership region of central North America.



"Barnyard Lignite," North Dakota, 1940



Indiana Steel, ca. 1910



Signal Hill, California, ca. 1923




Modern refinery, Billings, Montana, 2004

A Change Is in the Air

Just 200 years ago, before the onset of the Industrial Revolution, we had modest needs for energy and most of the energy we used came from what we now know as renewable sources. We used horses for transportation, burned animal dung and wood for heat, and used vegetable or animal oils in lamps for light. Water supplied much of the needed power to grind grain and for the limited manufacturing of the time. But as the Industrial Revolution


moved forward, largely on the shoulders of the steam engine, better energy sources were needed to fuel factories and transportation and provide energy to generate electricity. Energy-rich fossil fuels, including coal, oil, and natural gas worked well for these needs. Fossil fuel use has continued to increase dramatically in the industrialized world in the last 150 years.¹

Energy Consumption in the United States²



1. First commercial U.S. coal production begins near Richmond, Virginia³
2. James Watt patents modifications to steam engine³
3. Baltimore, Maryland, becomes first city to light streets with gas from coal³
4. First steamship to cross Atlantic³
5. Distillation of kerosene replaces whale oil⁴
6. First oil well in United States⁴
7. Edison invents electric lighting⁵
8. First commercial electric power station opens in San Francisco⁵
9. First practical coal-fired electric generating station goes into operation to supply household lights in New York.⁵
10. Steam turbine invented⁵

11. Gasoline-powered internal combustion engine developed³
12. Svante Arrhenius is first to investigate the effect that doubling atmospheric carbon dioxide would have on global climate.⁶
13. Electric refrigerator invented⁵
14. 9 million autos in the United States⁵
15. U.S. population at 148.7 million⁷
16. First commercial nuclear power plant⁵
17. 61.6 million autos registered in the United States⁸
18. Beginning of the modern global warming debate⁶
19. 129.7 million autos registered in the United States⁸ and an estimated 600 million motor vehicles in the world⁹
20. U.S. population at 281.4 million⁷



Greenhouse Effect


Energy from the sun drives the earth's weather and climate and heats the earth's surface; in turn, the earth radiates energy back into space. Certain atmospheric gases (water vapor, carbon dioxide, and other gases) trap some of the outgoing energy, retaining heat somewhat like the glass panels of a greenhouse.

Without this natural "greenhouse effect," global temperatures would be considerably lower than they are now, and life as it is known would not be possible. However, problems may arise when the atmospheric concentration of greenhouse gases increases.¹⁰




Nearly 100 years ago, Swedish scientist and Nobel Prize winner Svante Arrhenius postulated that anthropogenic increases in atmospheric CO₂ as the result of fossil fuel combustion would have a profound effect on the heat budget of the earth. In 1904, Arrhenius became concerned with rapid increases in anthropogenic carbon emissions and recognized that "the slight percentage of carbonic acid in the atmosphere may, by the advances of industry, be changed to a noticeable degree in the course of a few centuries."¹¹

Human (anthropogenic) activity, including the use of fossil fuel, generates a significant volume of greenhouse gases like CO₂. Since the beginning of the Industrial Revolution, atmospheric concentrations of carbon dioxide have increased nearly 30%, methane concentrations have more than doubled, and nitrous oxide concentrations have risen by about 15%.¹⁰ These increases have enhanced the heat-trapping capability of the earth's atmosphere. There is concern that the anthropogenic greenhouse gases entering the atmosphere are causing increased warming and that this warming will affect climate on a global scale.



Since the beginning of large-scale industrialization about 150 years ago, the level of CO₂ in the atmosphere has increased by about 30%.



The cycle of carbon movement between the biosphere, atmosphere, hydrosphere, and geosphere is a complex and important global cycle. In the atmosphere, carbon occurs primarily as carbon dioxide. Across the landscape, carbon occurs mainly in living organisms and decaying organic matter in soils.

Carbon is continuously circulated between reservoirs in the ocean, land, and atmosphere, where it occurs primarily as carbon dioxide. On land, carbon occurs primarily in living biota and decaying organic matter. In the ocean, the main form of carbon is dissolved carbon dioxide and small creatures, such as plankton. The largest reservoir is the deep ocean, which contains close to 40,000 GtC, compared to around 2000 GtC on land, 750 GtC in the atmosphere, and 1000 GtC in the upper ocean. Although natural transfers of carbon dioxide are approximately 20 times greater than those due to human activity, they are in near balance, with the magnitude of carbon sources closely matching those of the sinks.


What Is CO₂?


Carbon dioxide (chemical formula CO₂) is a colorless, naturally occurring gas composed of one atom of carbon and two atoms of oxygen. At temperatures below -78°C (-109°F), CO₂ condenses into a white solid called dry ice. When warmed, dry ice vaporizes directly from a solid to CO₂ gas in a process called sublimation. With enough added pressure, liquid carbon dioxide can be formed.

We use CO₂ to make the bubbles in soft drinks, and CO₂ (as dry ice) is used to keep things cold. Carbon dioxide is also used in fire extinguishers (CO₂ displaces the oxygen the fire needs to burn).

Presently, the earth's atmosphere is composed of about 0.04% CO₂. This small amount plays an important role in maintaining the natural greenhouse effect that makes Earth hospitable to life. Carbon dioxide is essential to plant life and is a key part of the global carbon cycle. An important component of this cycle takes place when plants take in CO₂ and break it down into carbon and oxygen through the process of photosynthesis. The carbon is stored in the plant and the oxygen is released to the atmosphere. When the plant dies or it is burned, much of the carbon recombines with oxygen to again form CO₂.

Carbon dioxide formed through human action is referred to as anthropogenic CO₂. Examples of activities that result in the production of anthropogenic CO₂ include plowing land, which exposes the carbon in the soil to the oxygen in the air; heating limestone to make lime for cement, which releases carbon in the limestone that combines with oxygen in the air; and burning fossil fuels for energy which combines the carbon in the fuel with oxygen. This anthropogenic CO₂ adds additional carbon to the global carbon cycle.






Carbon Management

To stabilize the levels of CO₂ in the environment, there needs to be a reduction in the amount of CO₂ that is released by human activity. Reducing anthropogenic CO₂ emissions with the goal of stabilizing the level of CO₂ in the atmosphere is called carbon management. Carbon management is a complex issue because most of the anthropogenic CO₂ comes from the use of fossil fuels for energy, and maintaining our energy flow is critical to our economy and our quality of life. Carbon dioxide emissions can be reduced by energy conservation, the use of more efficient fossil fuel energy systems, increased use of renewable and nuclear energy, and carbon sequestration.

Carbon sequestration, also known as carbon capture and storage (CCS), is the capture and storage of CO₂ and other greenhouse gases that would otherwise be emitted to the atmosphere and potentially contribute to global climate change. The greenhouse gases can be captured at the point of emission, or they can be removed from the air. Captured gases can be used; stored in underground reservoirs or, possibly, the deep oceans; absorbed by trees, grasses, soils, and algae; or converted to rocklike mineral carbonates or other products. Carbon sequestration holds the potential to substantially reduce greenhouse gas emissions.

There are two types of sequestration: *direct* and *indirect*. Direct, or geologic, sequestration involves capturing CO₂ at a source before it can be emitted to the atmosphere. The most efficient concept would use specialized processes to capture CO₂ at large stationary sources like factories or power plants and then inject the CO₂ into secure storage zones deep underground (geologic sequestration) or into the deep ocean. Indirect, or terrestrial, sequestration removes CO₂ from the atmosphere. Indirect sequestration employs land management practices that boost the ability of natural CO₂ sinks, like plants and soils, to remove carbon as CO₂ from the atmosphere, regardless of its source. Opportunities for indirect sequestration can be found in forests, grasslands, wetlands, and croplands.





Anthropogenic CO₂ can be captured and sent to storage before it enters the atmosphere (direct sequestration) or land management practices can be used to enhance carbon uptake after it enters the atmosphere (indirect sequestration).

Geologic Sequestration

Geologic sequestration involves the capture of anthropogenic CO₂ before it is released to the atmosphere and then injection into deep underground geologic formations. These formations, or CO₂ sinks, exist in a variety of configurations in sedimentary basins and include unminable coal beds, oil and gas reservoirs, and deep geologic layers that contain salty water (brine formations).

To be considered for sequestration, geologic sinks must have the characteristics that can hold the CO₂ in place for a long period of time (for example, a seal above a permeable zone of rock similar to the situation that would trap and store oil or natural gas), be isolated geologically from underground sources of drinking water, and be in a stable area (that is, an area not prone to earthquake activity). Successful geologic sequestration requires that the CO₂ stay in place and not pose a danger to human health and the environment.

Pure CO₂ has been stored as a gas in natural underground deposits for millions of years, and since the 1970s, oil field operators in west Texas have safely pumped millions of tons of CO₂ underground into oil-producing formations to increase production.

Under high-temperature and high-pressure conditions, such as those encountered in deep geologic formations (greater than 2600 feet), CO₂ will exist in a dense phase that is referred to as “supercritical.” When injected into a geologic formation, a portion of the supercritical CO₂ may be dissolved in any fluids, such as water or oil, that are present in the formation, while another portion will be available to react with rock minerals. When CO₂ dissolves in oil, it acts as a solvent, reducing oil viscosity and increasing its mobility. The sequestration of CO₂ in a supercritical form is beneficial for two reasons: 1) the supercritical state maximizes the number of CO₂ molecules that can be injected into a given volume and 2) if injected into an oil reservoir, supercritical CO₂ can increase oil production, which, in turn, can be used to pay for the capture and transportation of the CO₂ from the original source.

As with many disciplines and technologies, a precise and descriptive vocabulary is needed to adequately describe and discuss the sequestration of CO₂ in geologic formations. In the petroleum industry, a rock layer that contains fluid or gas is referred to as a *reservoir*. A rock layer that oil or gas cannot flow through is referred to as a *trap* or a *cap*. In hydrogeology, a rock layer that contains water is referred to as an *aquifer*. A rock layer that contains water with dissolved solids (salt) concentrations higher than drinking water standards is commonly known as a *saline aquifer* or a *brine formation*. A rock layer that water cannot flow through is referred to as an *aquitard* or a *confining bed*.

Carbon dioxide can be geologically sequestered in sedimentary basins by the following mechanisms: 1) stratigraphic and structural trapping in depleted oil and gas reservoirs, 2) solubility trapping in reservoir oil and formation waters, 3) adsorption trapping in unminable coal seams, 4) cavern trapping in salt structures, and 5) mineral immobilization.



NASA's Earth Observatory has created a series of images that show the color of the Earth's surface for each month of 2004 at very high resolution (500 meters/pixel) at a global scale. This image is of North America in June.

Terrestrial Sequestration


Terrestrial sequestration occurs at the Earth's surface through management practices that increase the amount of the carbon stored in roots and vegetable matter in the soil. It is important to remember that terrestrial sequestration does not store CO₂ as a gas but stores the carbon portion of the CO₂. If the soil is disturbed and the soil carbon comes in contact with oxygen in the air, the exposed soil carbon can combine with O₂ to form CO₂ gas and reenter the atmosphere, reducing the amount of carbon in storage.

Promising land and water management practices that lead to terrestrial sequestration of carbon include conservation tillage, reducing soil erosion, and minimizing soil disturbance; using buffer strips along waterways; enrolling land in conservation programs; restoring and better managing wetlands; eliminating summer fallow; using perennial grasses and winter cover crops; and fostering an increase in forests. Typically land management practices that store carbon have other benefits to the ecosystem including biodiversity, water filtration, increased soil health and fertility, and many others.

Soil can only take in and store a limited amount of carbon. On average, after a 50- to 100-year time frame, the soils will have reached equilibrium and not accept any more carbon. Once this "steady state" has been reached, the carbon will remain sequestered in the soil as long as the land is undisturbed and conservation land management practices are continued.

Terrestrial sequestration is important because it can be implemented immediately and can begin to reduce atmospheric CO₂ levels in the next few years. Using terrestrial sequestration now means we can get started on reducing CO₂ levels in the atmosphere while we adopt other carbon control measures.






DOE's Carbon Sequestration Regional Partnerships


If the decision is made that carbon sequestration must be implemented in the United States on a broad scale and in a relatively short time frame (meaning over the next 10–20 years), it will take a concerted effort of federal and state agencies working in cooperation with technology developers, regulators, and others to put into place both the economic framework and the necessary infrastructure to achieve meaningful carbon reductions.

To ensure that America is fully prepared to implement this climate change mitigation option, then-Secretary of Energy Spencer Abraham on November 21, 2002, announced plans to create a national network of public-private sector partnerships that would determine the most suitable technologies, regulations, and infrastructure needs for carbon capture, storage, and sequestration in different areas of the country. The Secretary called the partnership initiative “the centerpiece of our sequestration program.” The partnerships are a key part of President Bush’s Global Climate Change Initiative (GCCCI).

On August 16, 2003, following a competitive evaluation, Energy Secretary Spencer Abraham named seven teams, called Regional Carbon Sequestration Partnerships, to evaluate and promote the carbon sequestration technologies and infrastructure best suited to their unique regions. The original partnerships included leaders from more than 140 organizations spanning 33 states, three American Indian nations, and two Canadian provinces. By February 2007, the partnerships had expanded to include 350 organizations spanning 41 states, three American Indian nations, and four Canadian provinces.¹²

Today, the seven partnerships are developing the framework needed to validate and potentially deploy carbon sequestration technologies. They are evaluating numerous sequestration approaches that have emerged in the last few years to determine which are best suited for specific regions of the country. They are also identifying possible regulations and the necessary infrastructure requirements should our society determine that sequestration be deployed on a wide scale in the future.







The PCOR Partnership

The Plains CO₂ Reduction (PCOR) Partnership is a diverse group of public and private sector stakeholders working together to better understand the technical and economic feasibility of capturing and storing CO₂ emissions from stationary sources of CO₂ in the central interior of North America. The PCOR Partnership is managed by the Energy & Environmental Research Center (EERC) at the University of North Dakota and is one of seven regional partnerships funded by the U.S. Department of Energy's (DOE's) Regional Carbon Sequestration Partnership Program and a broad array of project sponsors.

The PCOR Partnership has assessed and prioritized the opportunities for sequestration in the region and identified and worked to resolve the technical, regulatory, and environmental barriers to the most promising sequestration opportunities. At the same time, the PCOR Partnership has informed policy makers and the public regarding CO₂ sources, sequestration strategies, and sequestration opportunities.

- Based on available data, the states and provinces within the PCOR Partnership region contributed roughly 13 % of the total CO₂ emissions from stationary sources in the United States and Canada.¹³
- Enhanced oil recovery (EOR), depleted oil and gas zones, deep saline reservoirs, and unminable coals in the PCOR Partnership region represent opportunities for direct (geologic) sequestration projects.
- The PCOR Partnership region is currently home to the Weyburn direct sequestration EOR demonstration project. There are four additional CO₂ sequestration field validation test sites in the PCOR Partnership region.
- Croplands, wetlands, rangelands, and forests in the PCOR Partnership region represent opportunities for indirect (terrestrial) sequestration projects.
- The PCOR Partnership region is currently home to several indirect sequestration research projects involving wetlands, cultivated land, prairie, and forest.



Phase II Partners		 Energy & Environmental Research Center® UNIVERSITY of North Dakota Grand Forks	 ADVANCED GEOTECHNOLOGY	 AIR PRODUCTS	 Alberta ENERGY	 EUB Alberta Energy and Utilities Board			
 AGS Alberta Geological Survey	 Apache CANADA LTD.	 BASIN ELECTRIC POWER COOPERATIVE A Touchstone Energy® Cooperative	 BlueSource	 BRITISH COLUMBIA Ministry of Energy and Mines	 O O Carbozyme Inc.	 CEED® Center for Energy and Economic Development	 Great Plains SYN FUELS Dakota Gasification Company	 Ducks Unlimited Canada CANADA'S CONSERVATION COMPANY	 DUCKS UNLIMITED
 Eagle Operating Inc.	 EICCO	 Encore Acquisition Company	 Environment Canada	 EXCELSIOR ENERGY	 Fischer Oil and Gas, Inc.	 GREAT NORTHERN	 GREAT RIVER ENERGY® A Touchstone Energy® Cooperative		
 HESS	 ILLINOIS AND GAS COMPACT COMMISSION	 DNR	 LIGNITE Energy COUNCIL	 MEG Energy Corp	 Melzer Consulting	 minnesota power	 Minnkota Power COOPERATIVE, INC. Your Touchstone Energy® Partner	 MISSOURI DEPARTMENT OF NATURAL RESOURCES	
 MISSOURI RIVER ENERGY SERVICES®	 MONTANA-DAKOTA UTILITIES CO. A Division of MDU Resources Group, Inc.	 Montana Department of ENVIRONMENTAL QUALITY	 NATIONAL COMMISSION ON ENERGY POLICY	 Natural Resources Canada	 Nexant	 NORTH AMERICAN COAL CORPORATION			
 North Dakota DEPARTMENT OF COMMERCE OFFICE OF RENEWABLE ENERGY AND ENERGY EFFICIENCY	 North Dakota Department of Health	 NORTH DAKOTA GEOLOGICAL SURVEY	 NORTH DAKOTA OIL AND GAS DIVISION	 Industrial Commission of North Dakota Lignite Research, Development and Marketing Program	 Industrial Commission of North Dakota Oil and Gas Research Council	 NORTH DAKOTA NATURAL RESOURCES TRUST	 North Dakota Petroleum Council	 NDSU NORTH DAKOTA STATE UNIVERSITY	 OTTER TAIL POWER COMPANY
 PTTC TECHNOLOGY CONNECTIONS	 Prairie Public Television	 Pratt & Whitney A United Technologies Company	 RAMGEN POWER SYSTEMS	 RPS Energy	 Saskatchewan Industry and Resources	 SaskPower	 Schlumberger		
 Shell Canada Limited	 Spectra Energy	 SUNCOR ENERGY	 DEPARTMENT OF ENERGY AND ENVIRONMENT	 NETL NATIONAL ENERGY TECHNOLOGY LABORATORY	 USGS science for a changing world	 UNIVERSITY OF ALBERTA	 WESTERN GOVERNORS' ASSOCIATION	 WESTMORELAND COAL COMPANY	 Xcel Energy™

PCOR Partnership Partners

The PCOR Partnership is a collaboration of more than 65 public and private sector stakeholders that have expertise in power generation, energy exploration and production, geology, engineering, the environment, agriculture, forestry, and economics. Our partners are the backbone of the PCOR Partnership and provide data, guidance, and practical experience with direct and indirect sequestration, including value-added projects.

PCOR Partnership List of Partners

- University of North Dakota Energy & Environmental Research Center (EERC)
- Advanced Geotechnology, a division of Hycal Energy Research Laboratories, Ltd.
- Air Products and Chemicals
- Alberta Department of Energy
- Alberta Energy and Utilities Board
- Alberta Geological Survey
- Apache Canada Ltd.
- Basin Electric Power Cooperative
- Blue Source, LLC
- British Columbia Ministry of Energy, Mines and Petroleum Resources
- Carbozyme, Inc.
- Center for Energy and Economic Development (CEED)
- Dakota Gasification Company
- Ducks Unlimited Canada
- Ducks Unlimited, Inc.
- Eagle Operating, Inc.
- Eastern Iowa Community College District
- Encore Acquisition Company
- Environment Canada
- Excelsior Energy Inc.
- Fischer Oil and Gas, Inc.
- Great Northern Power Development, LP
- Great River Energy
- Hess Corporation
- Interstate Oil and Gas Compact Commission
- Iowa Department of Natural Resources – Geological Survey
- Lignite Energy Council
- MEG Energy Corporation
- Melzer Consulting
- Minnesota Power
- Minnkota Power Cooperative, Inc.
- Missouri Department of Natural Resources
- Missouri River Energy Services
- Montana–Dakota Utilities Co.
- Montana Department of Environmental Quality
- National Commission on Energy Policy
- Natural Resources Canada
- Nexant, Inc.
- North American Coal Corporation
- North Dakota Department of Commerce Division of Community Services
- North Dakota Department of Health
- North Dakota Geological Survey
- North Dakota Industrial Commission Department of Mineral Resources, Oil and Gas Division
- North Dakota Industrial Commission Lignite Research, Development and Marketing Program
- North Dakota Industrial Commission Oil and Gas Research Council
- North Dakota Natural Resources Trust
- North Dakota Petroleum Council
- North Dakota State University
- Otter Tail Power Company
- Petroleum Technology Transfer Council
- Prairie Public Television
- Pratt & Whitney Rocketdyne, Inc.
- Ramgen Power Systems, Inc.
- RPS Energy
- Saskatchewan Industry and Resources
- SaskPower
- Schlumberger
- Shell Canada Energy
- Spectra Energy
- Suncor Energy Inc.
- U.S. Department of Energy
- U.S. Geological Survey Northern Prairie Wildlife Research Center
- University of Alberta
- Western Governors’ Association
- Westmoreland Coal Company
- Wisconsin Department of Agriculture, Trade and Consumer Protection
- Xcel Energy


Coal
Gasification Facility



Coal-Fired
Power Plant



Refinery



Ethanol Plant



Agricultural
Processing
Plant



Gas Processing
Plant



Cement Kiln

Anthropogenic Sources

The world now produces 25 billion metric tons of CO₂ each year from human activity (anthropogenic CO₂).¹⁴ Humans contribute this anthropogenic CO₂ to the atmosphere primarily through the use of fossil fuels in transportation, electrical generation, heating and cooling, and industrial activities. Additional anthropogenic CO₂ is also generated through agriculture.

Together the United States and Canada generate about a quarter of the world's anthropogenic CO₂. The United States produces about 5.7 billion metric tons (24%) of the global total, and Canada adds another 0.4 billion metric tons (nearly 2%). About 28% of Canada's anthropogenic CO₂ and 10% of the anthropogenic CO₂ generated in the United States is generated in the PCOR Partnership region.



The PCOR Partnership is focused mainly on finding practical ways to manage CO₂ from major stationary sources related to electricity generation; energy exploration and production activities; agricultural processing; fuel, chemicals, and ethanol production; and various manufacturing and industrial activities. These major stationary sources account for about 58% of anthropogenic CO₂ in the region. The PCOR Partnership is also working to implement terrestrial sequestration options that can capture and store CO₂ from the atmosphere.



Major Stationary CO₂ Sources in the PCOR Partnership Region

Annual CO₂ Output (tons)

- 25–500,000
- 500,000–5,000,000
- 5,000,000–10,000,000
- 10,000,000–15,000,000
- 15,000,000–20,000,000




CO₂ Sources

The PCOR Partnership project has identified, quantified, and categorized 1106 stationary CO₂ sources in the region. These stationary sources have a combined annual CO₂ output of nearly 549 million tons or 9.59 trillion cubic feet. And, although not a target source of CO₂ for direct sequestration, the transportation sector contributes nearly 223 million additional tons of CO₂ to the atmosphere every year.¹³

The annual output from the various stationary sources ranges from 10 million to 18 million tons for the larger coal-fired electric generation facilities, to under 5000 tons for industrial and agricultural processing facilities. In some cases, the distribution of the sources with the largest CO₂ output is coincident with the availability of fossil fuel resources, namely, coal, natural gas, and oil. This relationship is significant with respect to geologic sequestration opportunities. Many of the smaller sources are concentrated around more heavily industrialized metropolitan regions such as southeastern Minnesota, southeastern Wisconsin, and eastern Missouri.





Distribution of Annual CO₂ Output from Major Stationary Sources in the PCOR Partnership Region



Classification of Major Stationary CO₂ Sources in the PCOR Partnership Region

CO₂ Source Types

- Ag-Related Processing
- Electrical Utility
- Industrial
- Petroleum and Natural Gas



CO₂ Source Types


The geographic and socioeconomic diversity of the PCOR Partnership region is reflected in the diversity of the carbon dioxide sources found there. CO₂ is emitted from electricity generation; energy exploration and production activities; agricultural processing; fuel, chemicals, and ethanol production; and various manufacturing and industrial activities. The majority of the region's emissions come from just a few source types. About two-thirds of the CO₂ is emitted during electricity generation. Additional significant emissions come from industrial sources, petroleum refining and natural gas processing, ethanol production, and agricultural processing.


The emission profile (i.e., the percentage of CO₂ emissions from various source types) for the Canadian portion of the PCOR Partnership is virtually identical to that of Canada as a whole. When compared to the total U.S. CO₂ emissions, the states in the PCOR Partnership region emit relatively more CO₂ from electric utilities and less from industries and transportation.¹³

While the CO₂ emissions from the individual PCOR Partnership point sources are no different from similar sources located around North America, the wide range of source types within the PCOR Partnership region offers the opportunity to evaluate the capture, separation, and sequestration of CO₂ in many different scenarios.



Annual CO₂ Output by Major Source Categories for the States and Provinces in the PCOR Partnership Region





Geologic Framework

The same geologic framework that makes a large percentage of the PCOR Partnership region a significant producer of fossil fuels also creates prime opportunities for CO₂ sequestration. The western two-thirds of the region is underlain by great thicknesses of sedimentary rocks that span the entire stratigraphic record. The remainder of the region is underlain by Precambrian igneous and metamorphic rocks of the Canadian Shield.

The most extensive sequence of rocks in central North America is represented by the Cretaceous-aged marine sediments that were deposited in the former western interior seaway. This ancient sea extended from the Gulf of Mexico, across the center of North America, to the Arctic Ocean. The deeper portions of these strata and the underlying paleozoic-aged sediments offer tremendous capacity for sequestration.



Former extent of the Cretaceous-aged western interior seaway.

As the sea retreated from the continent, deltaic and marginal marine environments were established. The remains of these ecosystems are evident in the vast subbituminous coal and lignite reserves of Alberta, Wyoming, Montana, and North Dakota. The unminable portions of these deposits also provide opportunities for CO₂ sequestration.

In the millions of years since the seaway retreated, the central portion of the North American continent has been relatively stable. This tectonic stability is of prime importance with respect to safe and secure CO₂ sequestration in deep geologic formations.

	Age Units	YBP (Ma)	Potential Sequestration Targets
Cenozoic	Quaternary	1.8	
	Tertiary	66.5	Coal Seams
Mesozoic	Cretaceous	146	Coal Seams Saline Formations
	Jurassic	200	
	Triassic	251	
	Permian	299	Oil Fields Saline Formations
	Pennsylvanian	318	
Paleozoic	Mississippian	359	Oil Fields Saline Formations Oil Fields
	Devonian	416	Oil Fields
	Silurian	444	Saline Formations
	Ordovician	488	Oil Fields Oil Fields/Saline Fms
	Cambrian	542	Oil Fields Saline Formations
Proterozoic			
Precambrian			




Sedimentary Basins

Several relatively large and deep basins are found in the PCOR Partnership region, each with a sedimentary cover thousands of feet thick. The basins in the PCOR Partnership region have significant potential as geologic sinks for sequestering CO₂. Geologic sinks that may be suitable for long-term sequestration of CO₂ include both active and depleted petroleum reservoirs, deep saline formations, and coal seams, all of which are common in these basins.

While general information on the structural geology, lithostratigraphy, hydrostratigraphy, and petroleum geology of these basins is available, additional characterization data for specific geologic sinks will be necessary. Rocks that have been explored or developed for hydrocarbon recovery have been geologically characterized to a great extent, while non-hydrocarbon-bearing zones (such as saline formations) will require significantly more geologic investigation prior to large-scale sequestration.

Midcontinental Rift System

The PCOR Partnership region includes other areas besides the major petroleum-producing basins that are underlain by thick sequences of sedimentary rock. One of the largest and most notable of these areas is the Midcontinental Rift System, which stretches from eastern Nebraska across central Iowa and south-central Minnesota to the western portion of Lake Superior. The sedimentary rocks of Midcontinental Rift System may be viable locations for CO₂ sequestration. Because oil and gas have never been discovered in the Midcontinental Rift System, very few deep wells have been drilled in the area; therefore, little is known about the characteristics of these rocks. Continued regional characterization activities being conducted under Phase II of the PCOR Partnership will result in a better understanding of the potential for the sedimentary rocks of the Midcontinental Rift System to sequester large volumes of CO₂.





Oil and Gas Fields

The geology of carbon dioxide sequestration is analogous to the geology of petroleum exploration; the search for oil is the search for sequestered hydrocarbons. Oil fields have many characteristics that make them excellent target locations to store CO₂. Therefore, the geologic conditions that are conducive to hydrocarbon sequestration are also the conditions that are conducive to CO₂ sequestration. The three requirements for sequestering hydrocarbons are a hydrocarbon source, a suitable reservoir, and an impermeable trap. These requirements are the same as for sequestering CO₂, except that the source is artificial and the reservoir is referred to as a sink.

A single oil field can have multiple zones of accumulation that are commonly referred to as pools, although specific legal definitions of fields, pools, and reservoirs vary for each state or province. Once injected into an oil field, CO₂ may be sequestered in a pool through dissolution into the formation fluids (oil and/or water), as a buoyant supercritical-phase CO₂ plume at the top of the reservoir (depending on the location of the injection zone within the reservoir), and/or mineralized through geochemical reactions between the CO₂, formation waters, and formation rock matrix.

Oil is drawn from the many oil fields in the PCOR Partnership region from depths ranging from 2500 to 4000 feet for the shallower pools and up to 12,000 to 16,000 feet for the deepest pools.

Although oil was discovered in this region in the late 1800s, significant development and exploration did not begin until the late 1940s and early 1950s. The body of knowledge gained in the past 60 years of exploration and production of hydrocarbons in this region is a significant step toward understanding the mechanisms for secure sequestration of significant amounts of CO₂.



Selected Alberta Oil Fields

- 100s of fields
- Potential incremental oil = 2 billion bbl
- Total CO₂ needed for EOR = 9542 Bcf

Selected Saskatchewan Oil Fields

- 11 fields
- Potential incremental oil = 331 million bbl
- Total CO₂ needed for EOR = 2652 Bcf

Selected Montana Oil Fields

- Ten fields
- Potential incremental oil = 390 million bbl
- Total CO₂ needed for EOR = 3120 Bcf

Selected Wyoming Oil Fields

- 17 fields
- Total cumulative production = 1524 million bbl
- Potential incremental oil = 381 million bbl
- Total CO₂ needed for EOR = 3049 Bcf

Selected Manitoba Oil Fields

- Three fields
- Potential incremental oil = 39 million bbl
- Total CO₂ needed for EOR = 319 Bcf

Selected North Dakota Oil Fields

- 28 fields
- Potential incremental oil = 262 million bbl
- Total CO₂ needed for EOR = 2095 Bcf

Buffalo Field, South Dakota

- Portions of this field are currently undergoing tertiary recovery operations using air injection.
- CO₂-based EOR may be technically feasible.

Selected Nebraska Oil Fields

- Ten fields
- Total cumulative production = 100 million bbl
- Potential incremental oil = 25 million bbl
- Total CO₂ needed for EOR = 199 Bcf

Enhanced Oil Recovery Potential in the PCOR Partnership Region

 Oil Field

 Sedimentary Basin

0 100 200 300 600 Kilometers

0 100 200 400 Miles

Enhanced Oil Recovery

Most oil is extracted from the ground in three distinct phases: primary, secondary, and tertiary (or enhanced) recovery. Natural pressures within the reservoir drive oil into the well during primary recovery, and pumps bring the oil to the surface. Primary recovery typically produces roughly 12%–15% of a reservoir's original oil. An additional 15%–20% of the original oil can be extracted through secondary recovery processes which involve injecting water to displace the oil.¹⁵

Conventional primary and secondary recovery operations often leave two-thirds of the oil in the reservoir. In the United States, EOR methods have the potential to recover much of that remaining oil, which is estimated to be 200 billion barrels.¹⁵ However, oil recovery is challenging because the remaining oil is often located in regions of the reservoir that are difficult to access, and the oil is held in the pores by capillary pressure.


Reconnaissance-level CO₂ sequestration capacities were estimated for selected oil fields in the Williston, Powder River, Denver–Julesberg, and Alberta Basins. Two calculation methods were used, depending on the nature of the available reservoir characterization data for each field. The estimates were developed using reservoir characterization data that were obtained from the petroleum regulatory agencies and/or geological surveys from the oil-producing states and provinces of the PCOR Partnership region. Results of the estimates for the evaluated fields (using a volumetric method) in the four basins indicate a storage capacity of over 1.1 billion tons of CO₂.

Aside from non-market-based incentives, CO₂ sequestration in many geologic sinks is not generally economically viable under current market conditions. However, EOR miscible flooding is a proven, economically viable technology for CO₂ sequestration that can provide a bridge to future non-EOR-based geologic sequestration. For example, a portion of the revenue generated by CO₂ EOR activities can pay for the infrastructure necessary for future geologic sequestration in brine formations. It is expected that major oil fields subjected to this type of recovery process would retain a significant portion of the injected CO₂ (including the amount recycled during production) as a long-term storage solution.


Storage and Incremental Recovery Through EOR in Selected Fields

Basin	Cumulative Incremental Recovery (million bbl)	CO ₂ Sequestration Potential (Bcf)	CO ₂ Sequestration Potential (tons)
Williston	1023	8186	501,900,647
Powder River	381	3049	186,949,845
Denver–Julesberg	25	199	12,211,839
Alberta	2000	9542^c	545,000,000*

**Values for the Alberta Basin were determined using a different methodology than the other basins and, therefore, may not be directly comparable to the other estimates. They are included in the table to provide insight regarding the general magnitude of CO₂ flood-related sequestration capacity and potential incremental oil production in Alberta.*



Lower Cretaceous Saline System Analysis



Mississippian Madison Saline System Analysis




Distribution of selected saline systems evaluated in the PCOR Partnership region.
Limits of the evaluated areas are based on the extent of readily available data.


Saline Formations

Saline formations within the PCOR Partnership region have the potential to store vast quantities of anthropogenic carbon dioxide. Two saline systems, the Mississippian Madison and the Lower Cretaceous, have been evaluated for their regional continuity, hydrodynamic characteristics, fluid properties, and ultimate storage capacities using published data.

The lateral extent of these formations, the current understanding of their storage potential gained through injection well performance, and the geographic proximity to major CO₂ sources suggest they may be suitable sinks for future storage needs. For example, reconnaissance-level calculations on the Mississippian System in the Williston Basin and Powder River Basin suggest the potential to store upwards of 60 billion tons of CO₂ over the evaluated region, while the Cretaceous System has the potential to store over 160 billion tons.^{16,17}


Formation	Basin	Estimated CO ₂ Capacity (billion tons)
<i>Lower Cretaceous System</i>		
Newcastle Formation	Williston and Powder River	42
Viking Formation	Alberta	100
Maha Formation	Denver-Julesberg	19
<i>Mississippian System</i>		
Madison Formation	Williston and Powder River	60





Major Coal Basins in the PCOR Partnership Region

-  Coal Basin
-  Area of Current Assessment




Sequestration in Coal

Many coal seams throughout central North America are too deep or too thin to be mined economically. However, many of these coals have varying amounts of methane adsorbed onto pore surfaces, and wells can be drilled into the coal beds to recover this “coalbed methane” (CBM). In fact, CBM is the fastest growing source of natural gas in the United States and accounted for 7.2% of domestic production in 2003.¹⁸

As with oil reservoirs, the initial CBM recovery methods, dewatering and depressurization, can leave methane in the coal seam. Additional CBM recovery can be achieved by sweeping the coal bed with CO₂, which preferentially adsorbs onto the surface of the coal, displacing the methane. For the coals in the PCOR Partnership region, it is possible that up to thirteen molecules of carbon dioxide can be adsorbed for each molecule of methane released, thereby providing an excellent storage sink for CO₂.¹⁹ Just as with depleting oil reservoirs, unminable coal beds may be a good opportunity for CO₂ storage.

Three major coal horizons in the PCOR Partnership region have been characterized with respect to CO₂ sequestration: the Wyodak–Anderson bed in the Powder River Basin, the Harmon–Hanson interval in the Williston Basin, and the Ardley coal zone in the Alberta Basin. The total maximum CO₂ sequestration potential for all three coal horizons is approximately 8 billion tons.^{20–22}


In northeastern Wyoming, the CO₂ sequestration potential for the areas where the coal overburden thickness is > 1000 feet is 6.8 billion tons. The coal resources that underlie these deep areas could sequester all of the current annual CO₂ emissions from nearby power plants for about the next 150 years.²²



Land Cover Map of the PCOR Partnership Region

Land Cover Classification

	Deciduous Forest (4.8%)
	Evergreen Forest (20.9%)
	Mixed Forest (5.7%)
	Shrubland (9.4%)
	Temperate Grassland (18.1%)
	Polar Grassland (1.1%)
	Cropland (30.0%)
	Rock, Burnt Areas, Sparse Vegetation (3.6%)
	Urban and Built-up (0.2%)
	Water Bodies (6.0%)
	Snow and Ice (0.1%)
	Wetland (0.2%)



Terrestrial Sequestration

In contrast to direct sequestration deep within the earth, the concept of terrestrial sequestration focuses on a more passive mechanism of CO₂ storage in vegetation and soils within a few feet of the earth's surface. From the Central Lowlands forests and cropland in the southeastern portion of the region, through the expansive grasslands and croplands of the northern Great Plains, to the northern boreal forests of Canada, the PCOR Partnership region has a rich agrarian history founded on fertile soils. However, as central North America developed into the pattern of land use seen today, much of the original soil carbon was lost to the atmosphere. In this setting, the most promising potential to sequester carbon would be to convert marginal agricultural lands and degraded lands to grasslands, wetlands, and forests when favorable conditions exist.²³

Terrestrial sequestration methods that enhance carbon buildup in biomass and soils include adopting conservation tillage, reducing soil erosion, and minimizing soil disturbance; using buffer strips along waterways; enrolling land in conservation programs; restoring and better managing wetlands; eliminating summer fallow, using perennial grasses and winter cover crops; and fostering an increase in forests.^{23,24} Managing soils for increased carbon uptake will pull CO₂ from the atmosphere for a 50–100-year time frame, after which the soils will have reached a new equilibrium, i.e., a point at which the total amount of carbon in the soil does not change over time.²⁵ Once a steady state has been reached, the carbon will remain sequestered until the land management practices change or some other event occurs. The manipulation of soils and biomass for carbon sequestration has the advantage that it can be implemented immediately without the need for new technologies.






Prairie

Pothole

Region

The Prairie Pothole
Region of the PCOR
Partnership Region




Prairie Pothole Region

The PCOR Partnership region includes the Prairie Pothole Region, a major biogeographical region that encompasses approximately 347,000 mi² (222.4 million acres) and includes portions of Iowa, Minnesota, Montana, North Dakota, and South Dakota in the United States and Alberta, Saskatchewan, and Manitoba in Canada.²⁶ Formed by glacial events, this region historically was dominated by grasslands interspersed with shallow palustrine wetlands. Prior to European settlement, this region may have supported more than 48 million acres of wetlands, making it the largest wetland complex in North America.²⁷ However, fertile soils in this region resulted in the extensive loss of native wetlands as cultivated agriculture became the dominant land use. Because of oxidation of organic matter by cultivation, agriculture has depleted significant amounts of soil organic carbon (SOC) in wetlands.

The prairie potholes are an important element of the prairie ecosystem. This region accounts for up to 70% of the wild duck production in North America²⁷ and provides important breeding and migratory grounds for many types of wildlife. In addition to wildlife benefits, the prairie potholes provide many other ecological benefits, such as reduced erosion, improved water quality, flood and storm buffering, and recreational opportunities.

Recent work by Ducks Unlimited, Inc., and U.S. Geological Survey scientists for the PCOR Partnership demonstrated that restoration of previously farmed wetlands results in the rapid replenishment of SOC lost to cultivation at an average rate of 1.1 tons acre⁻¹ yr⁻¹.²⁸ The finding that restored prairie wetlands are important carbon sinks provides a unique and previously overlooked opportunity to store atmospheric carbon in the PCOR Partnership region.





CO₂-Rich Gas in
a Pinnacle
Reef Structure




Out of the Air – Into the Soil



CO₂ in an Unminable
Lignite Seam



CO₂ in a Deep Oil Reservoir



Field Validation Test Sites

Through the fall of 2009, the PCOR Partnership will be developing and conducting four CO₂ sequestration field validation tests: three that will store carbon dioxide as a gas in the deep subsurface and one that will store carbon in the near-surface soils and sediments of wetlands and grasslands. These field projects are designed to develop the expertise, real-world experience, and business models needed to implement major, full-scale, long-term CO₂ sequestration projects in the region.

CO₂-rich gas in a pinnacle reef structure – Acid gas (approximately 70% CO₂, 30% hydrogen sulfide [H₂S]) from a natural gas processing plant in northern Alberta, Canada, is being injected into an oil-producing zone in an underground pinnacle reef structure. A pinnacle reef is an isolated domed geologic formation suitable for trapping hydrocarbons and CO₂. Results will help to determine the best practices to support sequestration in these unique geologic structures as well as further our understanding of the effects of H₂S on tertiary oil recovery and CO₂ sequestration.


CO₂ in a deep oil reservoir – CO₂ will be injected into an oil-bearing zone at great depth in the Williston Basin in northwestern North Dakota. The activity will be used to determine the efficacy of CO₂ sequestration and the use of CO₂ to produce additional oil from deep carbonate source rocks.

CO₂ in an unminable lignite seam – CO₂ will be injected into unminable lignite seams in northwestern North Dakota. The injected CO₂ will be trapped by naturally bonding to the surfaces of the fractured lignite. The injected CO₂ also has the potential to displace methane occupying the coal fractures. This validation test will provide valuable information regarding lignites for both CO₂ sequestration and enhanced coalbed methane production.


Out of the air – into the soil – A managed wetland will be implemented in north-central South Dakota to demonstrate practices that will improve CO₂ uptake. The results will help to optimize CO₂ storage, monitoring, and verification methods and facilitate the monetization of terrestrial carbon offsets in the region and elsewhere.

Fossil Fuel Conversion Platforms and CO₂ Capture


Postcombustion



Precombustion



Oxygen
Combustion



Carbon Capture and Separation

Before CO₂ can be geologically sequestered, it must be captured and separated from other gases, compressed, and transported to the geologic sink. Capture and separation of CO₂ are performed as a part of many industrial processes, from H₂ production to ammonia synthesis.²⁹ However, existing capture technologies are not yet optimized for application on a typical power plant exhaust stream. Power plants typically burn coal or natural gas in the presence of air. This approach produces an exhaust gas that contains large amounts of nitrogen that must be separated from the CO₂ prior to sequestration. Because the concentrations of CO₂ in typical power plant flue gas are so low (ranging from 3% by volume for some natural gas-fired plants²⁹ to 15% by volume for some coal-fired plants³⁰), any capture process must be sized to handle the large quantity of exhaust gas. The large scale of equipment and quantities of chemicals required makes the capture process relatively expensive. In fact, the cost of capturing the CO₂ can represent three-fourths of the total cost of a capture-storage-transportation-geologic sequestration system.²⁹


Research is being performed to develop new CO₂ capture processes and improve the economics of existing ones. Capture of CO₂ can be performed at three points in the power production process: before combustion, during combustion, and after combustion. The precombustion technologies consist of capture of CO₂ in conjunction with gasification or reforming, while capture during combustion is only possible when the combustion takes place in nearly pure oxygen rather than air. The majority of capture technologies focus on separating the CO₂ from the exhaust gas stream after combustion.

Five different approaches can be taken in postcombustion capture: absorption (both chemical and physical), adsorption, membrane separation, cryogenic cooling, and others such as chemical looping and CO₂ hydrate formation. The most common commercial technology available is amine scrubbing, a chemical absorption technology. Two of the more common commercial physical absorption processes are the Rectisol® and Selexol™ processes.

After the CO₂ is captured, it must be compressed for either storage prior to truck transport or directly put into a pipeline to the sequestration site. CO₂ must be compressed to about 1200 to 1500 psi for transport in a pipeline. Compression is energy-intensive, so improved methods of compression are also being developed.



Top left and center: In the future, a more extensive CO₂ pipeline network will probably be constructed. Top right: Selexol™ towers at a recent integrated gasification combined-cycle start-up. Bottom: CO₂ compressors at the Dakota Gasification Company's synfuels plant in central North Dakota each move about 55 million cubic feet of CO₂ a day at a discharge pressure of 2700 psi.



Selected Technologies Used to Ensure the Safe and Secure Storage of CO₂

Terrestrial

Plant Matter Measurement

- Multispectral 3-dimensional aerial digital imagery
- Satellite imagery
- Light detection and ranging (LIDAR)

Soil Carbon Measurement

- Laser-induced breakdown spectroscopy (LIBS)
- Inelastic neutron scattering soil carbon

Geologic

CO₂ Fate and Transport Models

- Reservoir models (target formation to vadose)
- Geochemical models
- Geomechanical models
- CO₂ equation of state at reservoir conditions

Plume Tracking

- Surface to borehole seismic monitoring
- Microseismic monitoring
- Cross-well tomography
- Reservoir pressure monitoring
- Observation wells/fluid sampling

CO₂ Leak Detection

- Vadose zone soil/water sampling
- Air samples/gas chromatometry
- Infrared-based CO₂ in air detectors
- Vegetation growth rates
- CO₂ tracers, natural and introduced
- Subsurface monitoring wells

Mitigation

- Depressurize target formation

Regulatory and Safety Aspects of Carbon Capture and Storage

Currently, no U.S. federal regulations address CO₂ emissions. However, regulations do address the transportation of CO₂, the injection of CO₂ for enhanced resource recovery purposes, and worker safety issues. Also, various state and federal regulations could affect a CO₂ sequestration project. Most of these regulations would have bearing on the siting of a potential CO₂ source (e.g., power plant), pipeline routing, and injection of the CO₂. Further, numerous federal, state, and regional regulatory and/or legislative actions are being contemplated that would address various aspects of carbon management. Additionally, in April 2007, the Supreme Court decided that the U.S. Environmental Protection Agency (EPA) has the authority to regulate carbon dioxide emissions from cars and that the agency cannot bypass its authority to regulate greenhouse gases that contribute to global climate change unless it provides a scientific basis for its refusal. For more information on this Supreme Court decision go to www.supremecourtus.gov/opinions/06pdf/05-1120.pdf.

To ensure the safe and effective terrestrial and geological storage of CO₂, projects must identify and evaluate potential ecological and environmental impacts, effectively monitor and assess storage efficiency, and be prepared to take remedial action in the event of failure.

Assessing the effectiveness of terrestrial or geologic sinks for storing CO₂ is critical. Monitoring, mitigation, and verification (MMV) strategies will be required through all phases of CO₂ sequestration, including capture and separation, transportation, injection, and long-term storage.

The implementation of MMV serves to 1) protect worker health and safety; 2) ensure environmental and ecological safety; 3) verify safe and effective storage, including providing assurances of carbon credits of transactions in a carbon-trading market; 4) track plume migration; 5) provide early warning for failure; and 6) confirm model predictions.



What is known to date concerning risks of CO₂ sequestration includes the following:

- CO₂ can be safely stored in geological formations over long periods of time as observed with naturally existing CO₂ reservoirs.
- Environmental and ecological health effects are well understood.
- The largest risks of CO₂ capture and storage have been identified.
- Local hazards are generally more dependent on the nature of the release than the size of the release.
- CO₂ poses no health and safety risk at low concentrations.
- CO₂ is not flammable or explosive but does react with water.
- CO₂ is denser than air and has the potential to pool in low-lying areas or poorly ventilated spaces.



Decision Support System

PCOR Partnership

PCORP GIS

Indirect Carbon Sequestration

Regulating CO₂ Sequestration

Partner Contacts

- Phase II
- Phase I

Presentations

Products/Reports

- Phase II
- Phase I

DOE Global Climate Change Initiative Links

Other Carbon Related Links

- Sequestration / Storage / Infrastructure
- Data
- General/Background Information
- CO₂ Related Projects

Zama Zone

Terrestrial Zone

Contact Us

PCORP Home

Sponsored in part by the U.S. Department of Energy

© 2007 Energy & Environmental Research Center

The Decision Support System was prepared with the support of the U.S. Department of Energy Award No. DE-FC26-02NT41902. However, any opinions, findings, conclusions herein are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Energy.

DSS Decision Support System



Click on a site to view field validation test information.
Click [here](#) to view PCOR Partnership monthly updates.

The PCOR Partnership Decision Support System (DSS) is a web-based database, GIS, and scenario modeling system. The DSS allows our research partners to browse, query, analyze, and download data regarding CO₂ sequestration in the **Plains CO₂ Reduction Partnership**. This system is to compile an interactive interface that will provide for the defining range of transport and sequestration.

PCOR Partnership

Plains CO₂ Reduction (PCOR) Partnership

Practical, Environmentally Sound CO₂ Sequestration

EERC

University of North Dakota

Oil Field: Antelope

Reference: [NDIC](#)


Surface Area: 15,751 acres
Field Status: Non-Utilized
State/Prov: ND

	Pools	Unit	100%	75%	50%
Total Est. CO ₂ Capacity (mass)	9/9	tons	201,937,323	151,452,992	100,968,662
Total Est. CO ₂ Capacity (volume)	9/9	MMcf	3,293,598	2,470,198	1,646,799
Total Est. EOR CO ₂ Capacity (mass)	6/9	tons	47,421,456	35,566,092	23,710,728
Total Est. EOR CO ₂ Capacity (volume)	6/9	MMcf	773,444	580,083	386,722

[CO₂ Sequestration Calculations](#)

	Date	Oil (barrels)	Gas (Mcf)	Water (barrels)
Cumulative Production	3/2004	40,604,606	53,253,898	43,672,109

POOLS



The PCOR Partnership Decision Support System (DSS)


The PCOR Partnership has accumulated a wealth of data in characterizing the partnership region with respect to CO₂ sequestration opportunities. Major components of this characterization include creating an inventory of large stationary sources of CO₂ and identifying and mapping geologic and terrestrial targets, or sinks, for CO₂ sequestration across the PCOR Partnership region. Knowledge of the character and spatial relationships of sources, sinks, and regional infrastructure is crucial to developing and assessing approaches to economical and environmentally sound CO₂ sequestration.

The most efficient way to communicate this information to the partners has been through a geographic information system (GIS)-enabled Web site. This site is a major component of a larger Web-based Decision Support System® (DSS) that provides the PCOR Partnership with a single point of access to a wide variety of research data for evaluation and the development of potential sequestration scenarios. This password-protected (members only) Web-based platform contains the tools and capabilities designed to deliver functional and dynamic access to data acquired through the project. The data are housed in a relational database and accessed through a map-based portion of the Web site. More traditional Web pages provide access to relatively static data, such as reports, CO₂-related Web sites, terrestrial maps, and snapshots of regional data.

GIS technology enhances the users' understanding of regional opportunities by allowing them to visualize the spatially distributed nature of the data. The Web-based GIS interface contains several analytical methods that allow members of the research teams to browse, query, analyze, and download data regarding CO₂ generation and sequestration in the PCOR Partnership region. Researchers can use the GIS to:

- Examine attributes of individual features or groups of features and their spatial relationships to other features.
- Query the underlying data to analyze the region and export selected data for manipulation in other software.
- Explore the nature of the data through thematic maps.






Keeping the Lights On

Affordable energy not only fuels our vehicles and electrical plants, it also fuels our economy and our quality of life. Collectively, the states and provinces of the PCOR Partnership region use approximately 1200 trillion Btu of energy a year.^{31,32} At the most basic level, energy is essential, but to use our resources in a sensible way without damaging our planet requires a balance between energy and the environment.

The abundant, affordable energy provided by the PCOR Partnership region's fossil fuel resources powers a very productive part of the world. For example, the three Canadian provinces of the PCOR Partnership produce over 90% of Canada's wheat, while the U.S. portion of the PCOR Partnership contributes over 30% of U.S. wheat production.³³ Most of the continent's barley crop, which is critical to the breweries of Milwaukee and Saint Louis, comes from North Dakota and Minnesota. Wisconsin, as the top producer of paper in the United States, generates over \$12 billion in annual shipments of paper products.³⁴ The Missouri and Mississippi Rivers, railways, and highways of the region transport industrial output which includes heavy machinery, construction materials, and many other consumer goods.

The PCOR Partnership is working to develop technologies that will allow for CO₂ capture and sequestration. It is critical that technologies to reduce the environmental effects of fossil fuel use continue to be evaluated and developed while we explore and further develop future energy sources. The wise stewardship of our technological, social, and natural resources is essential to the future of our culture. Our challenge is to keep the lights on while simultaneously ensuring that our environment and economy stay strong.

Annual Energy Consumption



- 1 Sequestration PowerPoint presentations geared to general audiences, educators, students, and decision makers
- 2 PCOR Partnership’s public access Web site with sequestration background, streaming video, educational resources, and information links
- 3 PCOR Partnership’s Members-Only Web site with Decision Support System and Sequestration Project Resources
- 4 PCOR Partnership’s Technical Reports on a variety of sequestration topics in the PCOR Partnership region
- 5 A Sequestration Atlas for the PCOR Partnership region designed to inform the general public, educators, and decision makers
- 6 Fact sheets that summarize key sequestration topics and projects in the PCOR Partnership region
- 7 30-minute, broadcast-quality DVDs, coproduced by Prairie Public Television and the PCOR Partnership, on key sequestration topics
- 8 Proceedings of the PCOR Partnership’s annual meetings featuring PowerPoint presentations and program updates
- 9 PCOR Partnership’s display booth and tailored outreach packets available for informational meetings and educational venues



Education and Outreach – CO₂ Sequestration

The PCOR Partnership fully recognizes that the changes in land management, industrial operations, and CO₂ transport infrastructure needed to make large-scale, practical, and environmentally sound CO₂ sequestration a reality in the region cannot occur without an informed and supportive public.

For this reason, the PCOR Partnership has developed a number of outreach tools intended to educate and inform the public and decision makers about issues related to CO₂ and sequestration:

- A variety of PowerPoint presentations
- Display booth and materials
- Public Web site
- Members-only Web site
- Knowledge in brief – fact sheets on key topics and validation projects
- Knowledge in depth – scientific and technical reports
- Documentaries, available on DVD—coproductions of Prairie Public Broadcasting and the PCOR Partnership
- Proceedings from the annual PCOR Partnership meetings and access to other meeting materials

The PCOR Partnership also may have speakers for your meeting, school, or community group.

For more information regarding our outreach program or the content of this atlas and the Plains CO₂ Reduction Partnership, contact:

Edward N. Steadman
Senior Research Advisor
(701) 777-5279
esteadman@undeerc.org

John A. Harju
Associate Director for Research
(701) 777-5157
jharju@undeerc.org

Or visit our Web site at www.undeerc.org/pcor.

More information concerning DOE NETL's Regional Carbon Sequestration Partnerships can be found at www.netl.doe.gov/technologies/carbon_seq/partnerships/partnerships.html.



References

1. www.eia.doe.gov/oiaf/1605/ggccebro/chapter1.html (accessed August 2005).
2. www.eia.doe.gov/emeu/aer/contents.html (accessed August 2005).
3. http://instituteforenergyresearch.org/Energy_Timeline.htm (accessed August 2005).
4. www.sjgs.com/history.html (accessed August 2005).
5. www.eia.doe.gov/kids/history/timelines/index.html (accessed September 2005).
6. www.pbs.org/now/science/climatechange.html (accessed August 2005).
7. www.eia.doe.gov/emeu/aer/txt/ptb1601.html (accessed August 2005).
8. www.bts.gov/publications/national_transportation_statistics/2005/html/table_automobile_profile.html (accessed September 2005).
9. <http://hypertextbook.com/facts/2001/MarinaStasenko.shtml> (accessed August 2005).
10. <http://yosemite.epa.gov/oar/globalwarming.nsf/content/Climate.html> (accessed August 2005).
11. <http://earthobservatory.nasa.gov/Library/Giants/Arrhenius> (accessed August 2005).
12. www.fe.doe.gov/programs/sequestration/partnerships (accessed March 2007).
13. Jensen, M.D., Steadman, E.N., and Harju, J.A., 2005, CO₂ source characterization of the PCOR Partnership region: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005.
14. www.eia.doe.gov/oiaf/1605/ggccebro/chapter1.html (accessed August 2005)
15. www.netl.doe.gov/scngo/Petroleum/publications/eordrawings/Color/colop.pdf (accessed September 2005).
16. Fischer, D.W., Smith, S.A., Peck, W.D., LeFever, J.A., LeFever, R.D., Helms, L.D., Sorensen, J.A., Steadman, E.N., and Harju, J.A., 2005, Sequestration potential of the Madison of the northern Great Plains aquifer system (Madison geological sequestration unit): Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005.
17. Fischer, D.W., Sorensen, J.A., Smith, S.A., Steadman, E.N., and Harju, J.A., 2005, Potential CO₂ storage capacity of the Lower Cretaceous aquifer system in the PCOR Partnership area: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, September 2005.
18. www.fe.doe.gov/programs/oilgas/environment/Water_Resources_R&D.html (accessed September 2005).
19. Burruss, R., 2003, CO₂ adsorption in coals as a function of rank and composition—a task in USGS research on geologic sequestration of CO₂: The Second International Forum on Geologic Sequestration of CO₂ in Deep, Unmineable Coalseams (Coal-Seq II), March 6 and 7, 2003, in Washington, D.C., www.coal-seq.com/Proceedings2003/Burruss.pdf.
20. Nelson, C.R., Steadman, E.N., and Harju, J.A., 2005, Geologic CO₂ sequestration potential of lignite coal in the U.S. portion of the Williston Basin: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005.
21. Bachu, S.B., Steadman, E.N., and Harju, J.A., 2005, Carbon dioxide storage capacity in Upper Cretaceous–Tertiary Ardley coals in Alberta: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005.
22. Nelson, C.R., Steadman, E.N., and Harju, J.A., 2005, Geologic CO₂ sequestration potential of the Wyodak–Anderson coal zone in the Powder River Basin: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005
23. Paustian, K.H., and Cole, C.V., 1998, CO₂ mitigation by agriculture—an overview: Climatic Change, v. 40, p. 135–162, Netherlands, Kluwer Academic Publishers.
24. Peterson, E.B., Bonnor, G.M., Robinson, G.C., and Peterson, N.M., 1999, Carbon sequestration aspects of an afforestation program in Canada’s prairie provinces: Submitted to Joint Forest Sector Table/Sinks Table, National Climate Change Process, published by Nawitka Renewable Resource Consultants Ltd., p. 1–98, www.nccp.ca/NCCP/pdf/Afforest_Prairies.pdf (accessed September 2004).
25. de Silva, L.L., Cihacek, L.J., Leistritz, F.L., Faller, T.C., Bangsund, D.A., Sorensen, J.A., Steadman, E.N., and Harju, J.A., 2005, The contribution of soils to carbon sequestration: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005.
26. Mann, G.E., 1974, The Prairie Pothole Region—zone of environmental opportunity: Naturalist, v. 25, no. 4, p. 2 C. 7.
27. Wilkins, K.A., Otto, M.C., and Koneff, M.D., 2006, Trends in duck breeding populations, 1995–2006: U.S. Department of the Interior, Washington, D.C., 26 p.
28. Gleason, R.A., Euliss, N.H., Jr., McDougal, R., Kermes, K.E., Steadman, E.N., and Harju, J.A., 2005, Potential of restored prairie wetlands in the glaciated North American prairie to sequester atmospheric carbon: Plains CO₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, July 2005.
29. <http://fossil.energy.gov/programs/sequestration/capture/index.html> (accessed February 2007).
30. www.netl.doe.gov/technologies/carbon_seq/core_rd/co2capture.html (accessed February 2007).
31. www.statcan.ca/english/research/11-621-MIE/11-621-MIE2005023.htm#2 (accessed August 2005).
32. www.eia.doe.gov/emeu/states/_use_multistate.html (accessed August 2005).
33. <http://wbc.agr.state.mt.us/prodfacts/maps/wwwna.html> (accessed September 2005).
34. www.wipapercouncil.org/industry.htm (accessed September 2005).

Photo and Image Credits

Except for those noted below, all photos and images created by the EERC.

V.	Dragline with coal-fired electric generation station: Rick Craig Gas processing plant: Wolfgang Schlegl
Page 2.	Barnyard lignite: Vachon, John, 1940, FSA supervisor looking at pile of “barnyard lignite,” America from the Great Depression to World War II: Photographs from the FSA-OWI, 1935–1945. http://memory.loc.gov/ammem/fsowhome.html [fsa 8c18205] (accessed July 2005) Gary, Indiana Steel Plant: Haines Photo Company, 1910, Indiana Steel Co. #1, Gary, Indiana, Taking the Long View: Panoramic Photographs, 1851–1991. http://memory.loc.gov/ammem/collections/panoramic_photo/index.html [pan 6a19311 http://hdl.loc.gov/loc.pnp/pan.6a19311] (accessed July 2005) Signal Hill, California: Aerograph Company, 1923, Signal Hill, Taking the Long View: Panoramic Photographs, 1851–1991. http://memory.loc.gov/ammem/collections/panoramic_photo/index.html [pan 6a17401 http://hdl.loc.gov/loc.pnp/pan.6a17401] (accessed July 2005)
Page 4.	Pie chart data: www.eia.doe.gov/oiaf/1605/ggccebro/chapter1.html (accessed July 2004)
Page 5.	Image of Svante: http://cw.x.prenhall.com/petrucci/medialib/media_portfolio/text_images/FG14_19_05UN.jpg Earth from space: http://rsd.gsfc.nasa.gov/rsd/bluemarble/bluemarble2000.html (accessed September 2005)
Page 6.	Image of global carbon cycle: www.pmel.noaa.gov/co2/gif/globcar.png
Page 8.	Compact fluorescent bulb: Tiago Silva; Insulation of home: C.S. Poole; Solar panels: www.flickr.com (accessed May 2007) ; Wind turbines: Sandia National Laboratories
Page 9.	Weyburn photo: Courtesy of PTRC and the IEA GHG Weyburn CO ₂ Monitoring and Storage Project Farm photo: Photo by Lynn Betts, USDA Natural Resources Conservation Service
Page 12.	Tractor: Photo by Tim McCabe, USDA Natural Resources Conservation Service; Wetland: Photo by Lynn Betts, USDA Natural Resources Conservation Service; Grassland: Photo by Lynn Betts, USDA Natural Resources Conservation Service; Tree farm: USDA Natural Resources Conservation Service; Woodland: USDA Natural Resources Conservation Service
Page 13.	Norm Klopfenstein, USDA Natural Resources Conservation Service
Page 15.	U.S. DOE Carbon Sequestration Technology Development: U.S. Department of Energy, National Energy Technology Laboratory, 2007.
Page 17.	Harvest: www.deere.com/en_AU/images/press_releases/high_resolution_images/60serieshigh_2.jpg (accessed September 2005) Wind tower: www.sandia.gov/media/NewsRel/NR2000/wind.htm (accessed September 2005) Wetland: www.personal.psu.edu/users/m/e/mes270/Pictures (accessed September 2005) Pumpjack: www.copeland-corp.com/cp_ng/Images/Oil_Pump_Sunset.jpg (accessed September 2005) Dragline: www.lignite.com/mines/mines/Freedom%20Mine.htm (accessed September 2005)
Page 20.	Aerial of coal-fired electric generation station: Ed Murphy; Dragline and coal-fired electric generation station: Rick Craig; Gas processing plant: Wolfgang Schlegl; Ethanol plant: Lincolnland Agri-Energy; Cement kiln: Gas Technology Institute
Page 21.	Rush hour: www.raqc.org/images/RushHour.JPG ; Fireplace: www.stcloudheating.com ; Truck: www.qlinetrucking.com ; Train: www.arg.net.au/gallery.aspx ; Leafblower: Lukas Chatzioannidis; Mower: Gail Dedrick; Homes: Photo by Lynn Betts, USDA Natural Resources Conservation Service; Grill: www.flickr.com/photos/babygiu/195440147
Page 26.	Geologic map of North America: Kate E. Barton, David G. Howell, José F. Vigil, The North America Tapestry of Time and Terrain. U.S. Geological Survey Geologic Investigations Series I-2781 [http://pubs.usgs.gov/imap/i2781]
Page 27.	Evolution of life “Tower of Time” from John Gurche.
Page 29.	Cross section modified from Ground Water Atlas of the United States, Montana, North Dakota, South Dakota, Wyoming. HA 730-I. http://capp.water.usgs.gov/gwa/ch_i/gif/I011.gif (accessed August 2005)
Page 31.	Aerial photo: Bob Webster
Page 33.	EOR diagram modified from U.S. Department of Energy, National Energy Technology Laboratory, 1986. www.netl.doe.gov/scngo/Petroleum/publications/eordrawings/Color/colcarbon.pdf (accessed September 2005)
Page 34.	Mississippian Madison System: Fischer, D.W., Smith, S.A., Peck, W.D., LeFever, J.A., LeFever, R.D., Helms, L.D., Sorensen, J.A., Steadman, E.N., and Harju, J.A., 2005, Sequestration potential of the Madison of the northern Great Plains aquifer system (Madison geological sequestration unit): Plains CO ₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, June 2005. Lower Cretaceous System: Fischer, D.W., Sorensen, J.A., Smith, S.A., Steadman, E.N., and Harju, J.A., 2005, Potential CO ₂ storage capacity of the Lower Cretaceous aquifer system in the PCOR Partnership area: Plains CO ₂ Reduction (PCOR) Partnership Topical Report for U.S. Department of Energy and multiclients, Grand Forks, North Dakota, Energy & Environmental Research Center, September 2005.
Page 35.	Cross section modified from Ground Water Atlas of the United States, Montana, North Dakota, South Dakota, Wyoming. HA 730-I. http://capp.water.usgs.gov/gwa/ch_i/gif/I052.GIF (accessed August 2005)
Page 38.	Global Land Cover 2000 database. European Commission, Joint Research Centre, 2003, www.gvm.jrc.it/glc2000 (accessed June 2005)
Page 39.	Aerial view: Photo by Erwin Cole, USDA Natural Resources Conservation Service Grasses: Photo by Lynn Betts, USDA Natural Resources Conservation Service
Page 41.	Top: Prairie Pothole Region: Ducks Unlimited; Bottom: www.personal.psu.edu/users/m/e/mes270/Pictures (accessed September 2005).
Page 42.	Circle with wetland: Photo by Lynn Betts, USDA Natural Resources Conservation Service Circle with shovel of soil: Photo by Lynn Betts, USDA Natural Resources Conservation Service Lignite drilling: http://timberlineresources.biz
Page 45.	Top left: © BP p.l.c.; Top right: www.uop.com/objects/97%20Selexol.pdf ; Middle: http://library.enbridge.com Bottom left and bottom right: Great Plains Gasification Plant, Dakota Gasification Company
Page 47.	Gavel: www.justice.gov/gu (accessed December 2006)
Page 51.	Light bulb: http://gallery.hd.org/_c/light/_more2003/_more05/light-bulb-glowing-filament-light-blue-uncropped-lores-3-AHD.jpg.html (accessed August 2005) City at night: Dejah Thoris, www.flickr.com
Back	Palairret, J., and Delarochette, L., 1765, A Map of North America. From Library of Congress, Map Collections: 1500–2004. http://hdl.loc.gov/loc.gmd/g3300.ar011501 .

A Much Earlier Map of the PCOR Partnership Region

